

## MRD CHECKLIST

PARTICULARS	YES	NO
- IP Number allocated to each Patient	/	
- Name, Age & Sex of Patient	/	
- General Admission Consent	/	
- Initial Assessment of Patient / Diagnosis	/	
- Nutritional Assessment by Consultant	/	
- Plan of care counter signed by the Consultant		
- Treatment Orders - Date, Time, Name & Sign.	/	
- Medication Order / Drug Chart - Date, Time, Name & Sign.	/	
- Vital Signs Chart (TPR Chart)		-
- Intake Output Chart		
- Drug Chart (Duly filled)	/	
- Anesthesia Consent - (8 thing) - Date, Time, Name & Sign. of both Patient & Anesthetist		
- Anesthesia Assessment Sheet		
- Surgery Consent - (8 things) - Date, Time, Name & Sign of both Patient & Surgeon		
- Surgery Notes - Post Operative Plan		
- Pain Scoring System		
- Blood Transfusion if done		
- High Risk Procedures		
- A copy of the Discharge Summary	7	





59/Malc/MHI202481729 11/01/2024/IPH2024000093

Dr.G. GNANAVELU





# Medway Hospitals

(A Unit of United Alliance Healthcare Pvt Ltd)  ADMISSION SLIP
Admitting Doctor: Dr. Grandolu Speciality: Cardio 10014
Advised Date & Time: 91110696
Provisional Diagnosis:
ACS-ROCENT AWMS (COPI).
Reason for Admission: Medical Management Surgical Management
Others (please specify details)
dmission Type: Day Care ER Ward
CU (Specify details)
Surgery / Procedure Name (if planned):
CAY
Blood Product Requirement: No Yes (Kindly specify details of components required in space below)
<u> </u>
Expected Duration of Stay: Our large
Expected Cost of Treatment (as per Financial Counseling Form):
Payer: Self Insurance Others:
Instructions to Nurse (if any):  Admission in P
Any other Instructions (if any):
Doctor's signature Name Dr. G. Gnanavelu MD, DN Regio No. Co.  Advisor & Mer tor  Chief Cardiologist 39469 7 125 9.47

Reg. No: 39469

For admission desk staff only:				
Room Category: General Ward Single Room Twin Sharing Deluxe Room Suite Room Others				
Admission intimation		Admission Ti		
Date 11   1   24	Time	Date 11 24	Time	
Source: OPD  BR Direct  To be filled only if Blood requirement specified by the Doctor:  Is Blood Reservation and Blood Bank clearance completed as advised: Yes No				
Front office Staff Signature	Name Da.	P Emp. No.	Date   Time   10'   4	
		·		

-



# Medvay Hospitals The way to better health (A Unit of United Allience Mealths.)

(A Unit of United Alliance Healthcare Pvt Ltd)



## Mr.SAMRUDDIN S M

59/Malc/MHI202481729 11/01/2024/IPH2024000093

Dr.G. GNANAVELU





MHI/HOSP/2022/129

**ADMISSION FORM** 

Marital Status	Full Address / 23/12 Calama: Calama: Calama:	/ Telephone Number
M	Hillath Nogor Nethali Nagari	9 979017176
Occupation R L	Full Address   23/13 Jslamal Coloner Rose Fullath Nagari Nethali Nagari Varniyam badi	
Referred from		al No. of Days
Dr. a.	a. 11/1/24 (11/1/24 @ 8 h)	<u>.</u>
UNIT	MLC Yes No If Yes AR No.:	
	FINAL DIAGNOSIS	ICD Code
	CAP - EVOLVED AWMI	7561
	MORERATE LY DYSFUNCTION	TJ0:1
	COPP.	544.9
	·	
DATE	OPERATION / PROCEDURES	ICPM Code
اعدارا	CORDNARY ANGIOGIRAM PONE	02.88
11/1/24 ON 11/1/24		
DATE	TYPE OF ANESTHESIA	
14/1/24	GENERAL SPINAL COCAL REGIONAL	EPIDURAL
	DISCHARGE STATUS	
☐ Cured		
☐ Jmproved	☐ Against Medical Advice	Expired > 48 hours
☐ Unchang	☐ Absconded	Post-Operative Death
- In	C. Alex	loy cas
Signature	Ag : > '	dical Records Officer

## **AUTHORISATION FOR TREATMENT I PAYMENT**

administer such drugs as may be neces	sary and to perform suc the diagnosis and treat	Paramedical, Staf f of the Hospital Investigate treat and h operation under anaesthesia or other wise as may be ment of my illness / patient
I hereby under take to settle all the bills f basis. In any case, I shall pay all the due	·	es related to me/the patient named overleaf on a periodic ged from the hospital.
	•	agreed above, I hereby authorise the hospital to transferent as deemed fit and proper by the hospital authorities.
	r theis attendants have	nd Regulations of the Hospital and that all cash, jewellery been removed to a place of safety / handed over to the egard to any loss.
I have read out and explained the conter	nts of the above to the S	ignatory in his vernacular .
சிகீச்சை, பணம் செலுத்துதல் முதலியவை 6		
இதன் மூலமாத நான் நிர்வாகம், மருத்துவம்,	தாதியா், ஏனைய மருத்து க்கு தேவைப்பட்	வ ஊழியர்கள் எனக்க / நோயாளி . <u>இவரை படிய</u> படி இது கடிய கூறு மருந்துகளை கொடுக்கவும். மயக்க
		லவுகளை கட்டத் தவறினால் என்னை நோயாளியை வேறொரு ஒப்புதலை எனது உறவினர்கள் மூலமாக பெற நான் அதிகாரம்
மருத்துவமனையின் பொது சட்ட தீட்டங்கள்	பற்றி தெரிவிக்கீப்பட்டிருக்	<del>க</del> ீறேன்.
•	•	ட்கள் யாவும் பாதுகாப்பான இடத்திற்கு மாறுபட்டுவிட்டன / அல்லு ன எனது/நோயாளியின் எந்தவித நஷ்டத்திற்கு பொறுப்பில்லை
மேற்குறிப்பிட்ட அனைத்தும் எனக்கு விவரிக்	கப்பட்ட பிறகுதான் கைவெ	பாப்பமிட்டேன்.
R. Go		· 10 10 15 10 00
செவிலியர் கையொ:்பம்	தேதி	எனது/உறவினர்/காப்பாளர் கையொப்பம்
Signature of Admitting Nurse	Date 11/1/24	Signature of the Patient / Relative / Gurdian
		wife
		உறவுமுறை


Nature of Relationship



discharge.









## **GENERAL CONSENT FOR ADMISSION**

i,	the Patient or Representative of patient have lease tick the correct option above and below)
	lease tick the correct option above and below) ☑ Read
_	☐ Head☐☐ Been explained this consent form in English, which I fully understand.
_	Deer explained this consent to min English, which halfy dideratario.
•	I give my full consent and authorization for admission and treatment at this hospital. The proposed treatment plan has been explained to me.
•	I consent and authorize the hospital, treating doctors, nursing, technical and paramedical staff to provide relevant care and to conduct diagnostic as deemed necessary by the treating doctor/team.
•	I also consent to use of assistants such as resident doctors, other doctors, nurses, and other healthcare workers by the hospital and treating doctor / team.
•	l consent for clinical consultation, admission, disclosure of information required for clinical management (under confidence), routine medical examination (physical examination, palpation, percussion, auscultation), routine lab and imaging investigations, general nursing care, diet and physiotherapy assessment and counselling.
•	I have been explained about the proposed care plan, expected result(s), possible outcome(s) and expected cost of treatment/ hospital stay.
•	I understand that the hospital will take due care of me / my patient but, that there is always a possibility of an unexpected complication(s) which may necessitate longer stay and / or use of intensive care services. In such cases, procedure different from those contemplated and other intervention(s) may sometimes be needed.
•	I declare that, I have and will inform the doctor of my medical history including previous illnesses, allergies, drug reaction(s), surgical procedure, relevant medical family history and all other facts relevant to my treatment. I shall not hold the hospital/ doctor responsible for any consequences which may arise due to non-disclosure of relevant information on my part.
•	I declare that I have been explained about my rights and responsibilities.
•	I have been made aware of the rules and regulations of the hospital including those related to security and I promise to abide by them.
•	I understand that in case of some unexpected event occurring during the course of my stay I may be suggested a transfer to another hospital / healthcare organization, as considered appropriate by my treating doctor.
•	I understand that, drugs, consumables and devices will be charged on an 'as actual' basis as per the hospital tariff. I have been informed and I understand that there can be usage of certain reprocessed items during the course of the treatment. I also understand that only full strips of medicines shall be issued and returned. I declare that I take full responsibility of settling the bill before leaving the hospital premises at the time of

- I further declare that I have been given an opportunity to ask question(s) related to my admission, care plan and proposed hospital stay, and that such questions have been answered to my satisfaction.
- I declare that I have received and fully understood the information provided in this consent form, that I have been given an opportunity to ask questions relating to my admission, care plan and proposed hospital stay, and that all my questions have been answered to my entire satisfaction and there are no misconceptions or false hopes in my mind. I further declare that all fields (of this form) requiring insertion or completion were filled in my presence at the time of my signing this form.
- I, the above-named Patient / named patient's representative, do further hereby declare that I am above 18 years of age as on the date of signing this form, mentally sound and am giving consent without any fear, threat or false misconception.

	Signature / Thumb Impression*	Name	Date	Time	
Patient	7669500	Samueldin	11/1/24	10:14	
Surrogate/Guardian (if applicable #)	(J) 0 5 1 6		11/1/24	10:14	
Reason for surrogate consent	Patient is unable to give consent i	pecause:			
Witness	(voie stra M	um Hai	11/1/24	10:14	
Interpreter (if applicable)					

<sup>\*</sup> Right Hand for Males & Left Hand for Females | # Only if Patient is a minor or unable to give consent





Every heart beat counts (A Unit of United Alliance Healthcare Pvt Ltd)

## DAY CARE DISCHARGE SUMMARY

IP No.

IPH2024000093

D.O.A

: 11/01/2024

**UHID** 

MHI202481729

D.O.P

: 11/01/2024

Name

. Mr. SAMRUDDIN. S.M

Room No. : RI.

Age / Gender

59 Years /MALE

Consultant

Dr. G. Gnanavelu, MD., DM., (cardio) FACC

D.O.D

: 11/01/2024

Chief Cardiologist

#### **DIAGNOSIS:**

CAD - EVOLVED AWMI

MODERATE LV DYSFUNCTION

COPD

PROCEDURE: CORONARY ANGIOGRAM DONE ON 11.01.2024 - OSTIAL LAD NEAR TOTAL

OCCLUSION.

#### **BRIEF HISTORY:**

Mr. Samruddin. S.M, 59 years old male, presented with complaints of chest pain for 1 week & palpitation associated with sweating. He was evaluated in ESIC hospital and advised for Coronary angiogram and referred to Medway Heart Institute on 11.01.2024 for which he has been admitted.

#### ON EXAMINATION:

HR: 86bpm:

BP: 117/80 mmHg;

SPO<sub>2</sub>: 97% in room air

CVS: S1S2+; RS: Clear:

CNS: NFND;

Abd: Soft

#### INVESTIGATIONS:

BLOOD: Hb- 14.4gm/dl, TWBC - 7450cells/cumm, PLT - 178000 cells/cumm, Urea - 15.12mg/dl, Creatinine – 0.56mg/dl, Sodium – 138mg/dl, Potassium – 4.12mg/dl, PT /INR – 10.9/0.9.

ECG: sinus rhythm, HR @ 98bpm. Evolved AWMI changes.

**ECHO**: RWMA(+), Mid septal, mid anteroseptal hypokinesia. Distal septal, distal lateral apical hypokinesia. 1/4 MR. Severe LV dysfunction. EF – 33%. No PE / clot. No PHT.

#9, 1st Main Road, United India Colony, Kodambakkam, Chennai - 600024. Tel : 044 - 4310 8959

@MedwayHospitals

(a) @medwayhospitals

@medway-hospitals

@medwayhospitals

94557 94557 1800 572 3003

**Medway Group of Hospitals** 

E-mail: info@medwayhospitals.com | Website: www.medwayhospitals.com | CIN: U74900TN2011PTC083665

Medway Centre of Excellence (Chennai)

Kodambakkam Mogappair 044-2473 4455 | 044-26530011 | 044-27426829 | 04146-242000 | 044-2473 4455 | 0884-2333367

Chengalpattu

Villupuram

Kumbakonam

Kakinada

**Heart Institute** 044 - 4310 8959 Institute of Pulmonology 044-2473 4451



UHID: MHI202481729



#### Every heart beat counts

(A Unit of United Alliance Healthcare Pvt Ltd)

## CORONARY ANGIOGRAM FINDINGS:

Right -dominant system; OSTIAL LAD NEAR TOTAL OCCLUSION.(reports enclosed)

ADVICE: IVUS GUIDED PTCA TO LAD (POBA TO DIAGONAL) VS CABG.

#### ADVICE MEDICATIONS:

ŠI.	NAME OF THE DRUGS WITH	DOSAGE	FRE	QUEN	CY _	ROUTE	RELATION	DURATION
NO	GENERIC NAME		M	A	N		SHIP WITH FOOD	
1	TAB. ASA (ASPIRIN)	75 MG	0	1	0	ORAL	AFTER FOOD	TO CONTINUE
2	TAB. CLOPILET (CLOPIDOGREL)	75 MG	1	0	1	ORAL	AFTER FOOD	TO CONTINUE
3	TAB. ATORVA (ATORVASTATIN)	20 MG	0	0	1	ORAL	AFTER FOOD	TO CONTINUE
4	TAB. NITROCONTIN (NITROGLYCERIN)	2.6 MG	1	0	1	ORAL	AFTER FOOD	TO CONTINUE
5	TAB. ENVAS (ENALAPRIL)	2.5 MG	1	0	1	ORAL	AFTER FOOD	TO CONTINUE
6	TAB. MET XL (METOPROLOL)	25 MG	1	0	0	ORAL	AFTER FOOD	TO CONTINUE
7	TAB. LASIX (FUROSEMIDE)	40 MG	1/2	0	0	ORAL	AFTER FOOD	TO CONTINUE
8	TAB. ALDACTONE (SPIRONOLACTONE)	25 MG	I	0	0	ORAL	AFTER FOOD	TO CONTINUE
9	TAB. PAN (PANTOPRAZOLE)	40 MG	1	0	0	ORAL	BEFORE FOOD	TO CONTINUE

DISCHARGE ADVICE			
DIET	LOW FAT DIET.		
PHYSICAL ACTIVITIES	AVOID STRENUOUS ACTIVITIES.		
REVIEW	REVIEW WITH CARDIOLOGIST IN ESIC HOSPITAL.		

To report:

If temp > 101 'F / Difficulty in breathing / chest pain / Giddiness/ palpitations.

In case of emergency Contact: Medway Hospitals @ 4310 8959.

Dr. G. Gnanavelu. MD., DM., (cardio) FACC Chief Cardiologist

Typed by: Ezhilarasi.

Dr. G. Gnanavelu MD, DM (cardio), FACC

Chief Cardiologist Reg. No: 39469

Content of the

discharge summary." #9, 1st Main Road, United India Colony, Kodambakkam, Chennai - 600024. Tel : 044 - 4310 8959

**★** @MedwayHospitals

(C) @medwayhospitals

@medway-hospitals

@medwayhospitals

94557 94557 1800 572 3003

**Medway Group of Hospitals** 

E-mail: info@medwayhospitals.com | Website: www.medwayhospitals.com | CIN: U74900TN2011PTC083665

Medway Centre of Excellence (Chennai)

Kodambakkam Mogappair 044-2473 4455 | 044-26530011 | 044-27426829 | 04146-242000 | 044-2473 4455 | 0884-2333367

Chengalpattu

Villupuram

Kumbakonam

Kakinada

**Heart Institute** 044 - 4310 8959 Institute of Pulmonology 044-2473 4451





59/Male/MHI202481729 11/01/2024/IPH2024000093

Dr.G. GNANAVELU





Every heart beat counts

## DAY CARE INITIAL ASSESSMENT FORM

	TAI OAKE MITTAE AGGEOGRENT I OKNI					
Date	Date: 11   24 Time of arrival: 10.15.					
	A (to be filled by Nurses					
<b>Vital</b> Respi	Vital Signs: Temp: 并与(°F)   Pulse / HR: 是 (beats/min)   BP: 17   80 (mmHg)  Respiration: 20 (breaths/min)   SpO <sub>2</sub> : 年 (%)   Height: 以 (cms)   Weight: 54式 (kgs)   BMI: 21   kg (m)					
_	Any Language Barrier: Yes Any Language Coordinator / Translator  Allergies: Yes Yes Yes, specify:					
Alcol Do ye	Psychosocial Assessment:  Alcohol Intake:   Yes No Substance Abuse:  Yes No Smoking:  Yes No Smoking:  Yes No No Yes No No Yes Yes No Yes No Yes Specify details:					
Pain Screening Pain: Yes No. If Yes, Score: C C C Pain Scale used: PIPPS (28 weeks to < 38 weeks) CRIES (38 weeks - 2 months)  FLACC Scale (2 months - 7 years) Wong-Baker FACES Pain Rating Scale (7 years - 12 years)  Numerical Rating Scale (Age more than 12 years)  Duration: Location: Burning Referred / Radiant Pain						
Nutritional Screening:  Last 3 months Appetite ☐ Increased ☐ Decreased ☐ No Change  Last 3 months Weight ☐ Increased ☐ Decreased ☐ Mo Change						
Fall Risk Screening for adults: All Risk  Age more than 65 years History of fall in last 3 months  Walks with assistance Any neurological problem  In case of 2 or more criteria met initiate detailed fall assessment and fall prevention protocol						
Fall Risk Screening (for pediatrics)  H/O fall in last 3 months Neurological problem (vertigo, seizure, etc) Deranged Mobility No Risk  In case of 2 or more criteria met initiate detailed fall assessment and fall prevention protocol						
	Signature	Name	Emp. No.	Date	Time	
Nurse	Auf	Louranya.	0158.	11/11/24	10,20	









Every heart beat counts

# The way to better Mr.SAMRUDDIN S M

59/Malc/MHI202481729 - 11/01/2024/IPH2024000093

PR	OG	RESS	ИО	TES

	Dr.G. GNANAVELU	PROGRESS NOTES				
DATE	448 184 800 800 800 800 800 800 800 800 800 8	NOTES				
1-016						
11/1/24	CAG: Post Radu	il to state opth.				
10		Cook: Reg Radial or shall option.  Bro groo				
\d\'	gro of t	gro of For				
	<b>V</b>					
	Plan. Peg to Loo					
		932				
11.11						
11/1	pells on miny (can	)				
12.25	Pi. Rinnimo					
<u> </u>	- 110 s normal					
	CARG - SVD of LAND					
	- PMW - PCA					
	-10 26 DR.	GG & EJI FOR FEMILIAN PLANTS				
	h					
_	Mand					
	Dr. Afrish Nelson Reg. No: 88434					
1400	from can be ordered	rlag				
	Augan					
	Dr. Anish Nelson Reg. No: 88434	-				







# Every heart h

59/Maic/MHI202481729 11/01/2024/IPH2024000093

## Dr.G. GNANAVELU

## Department of Dietetics

## NUTRITION ASSESSMENT AND CARE PLAN FORM

nosis:	767/F	PCS - ALCOTO	MICOPO	15E-3	3-61	v v v	. 2
161	cms	Weight:Kgs	Food allergies:	Yes/ Nor if yes, specif	fy		
ious Beliefs:	, J	Vegetarian / /	Non Vegeta	rian - , , s	*	] Eggetarian	☐ Jain
Prescription:	600	calones	, con fat,	LOW SO	Ot i	00000	luid nest
ŊĘĊŢĮVI	E GĻOBA	AL ASSESSMENT	(ADULTS)	and the second	, , .	A. c	) <del>,</del>
	(A) -	Patient's related Medical Histor		<u>., .,</u>	_, _ (	1	<del></del>
	1)	Weight Change (overall-change	in past 6 months)	* j = 10 \$		1 1/1	<del></del> -
				l'@3	1		
	·	No weight change/	cSX' . To a fine a	5-10%	11 1	10-15%	>15%
2)	Dietary Intake	Duration:					
		1	D 2 '	3	· -'	<u> </u>	. □5
	Oral	No change	Sub - optimal solid diet	Full liquid diet/ moderate overall decrease	1 1	Hypo - caloric liquid diet	Starvation
-	Enteral / Parenteral Nutrition	Adequate / Excessive	Sub - optimal	Inadequate		Typo - caloric feeds	Starvation
	<del></del>		L		*, ,	l	
3)	Gastrointestin	al Symptoms Ouration:	<u> </u>	, , , , , , , , , , , , , , , , , , ,	<u>'                                     </u>	<u> </u>	
	V	1	<del></del>	11			<del></del>
	_	No symptoms	Nausea	Vomiting / moderate GI / , 1 symptoms	· · · · · · · · · · · · · · · · · · ·	Diarrhoea	severe anorexia
4)	Functional Ca	spacity (Migritton related functional Impa	Irment) Duration:				
	0	10 i		□3. '		□ 4	□ s
		None /Improved	Difficulty with ambulation	Difficulty with normal activity		Light activity	Bed / chair - ridden with no or little activity
5)	Co - morbidity	Disease and its relationship to nutrition	L		, ,,	f	
		X(1	□ ·2	3.	) · •	4	_ s
	` ` ` `	Healthy	"Mild co- morbidity	Moderate co- morbidity/ age >75 years		severe co - morbidity	Very severe multiple co - morbidity
B) ,	Physical exan	nination	I		· · ·	<del>                                     </del>	
1]	<del></del> -	stores or loss of subcutaneous fat			-	<u> </u>	
	Jed case in	1	□ 2,		· · · · · ·	. 4′ ,	<u> </u>
	<del>                                     </del>	Normal	Mild	Moderate	, , ,	<del></del>	Severe
2)	Sign of muscle v		ward	moderate ",		<del></del>	acres .
ㅋ	Tiller of thirty Clay	1			<u> </u>	□4.	
	+	Normal	Mild	Moderate		<del>  "</del> -	. Severe
Total Score = :	Sum fabove 7 comp	<u> </u>		Y		7- 1-	
Nutritional St	atus : Based on this						
	Well Nourished			(7 to 14)	ì	<del></del>	<del></del>
	Moderately Mai	<del></del>	<del></del>	(15 to 18)	-		
	Severely Malnor		<del>``</del>	(19 to 35)	-		
Nutrition Inte	ervention:					<del>.</del>	<del></del>
	Orai		lc	Enteral	☐ Paren	teral	· · · · · · · · · · · · · · · · · · ·
Diet counselli		□ Yes		No			<del></del>
	re-assessment:	Weekly		☐ Fort - nig	tht	☐ Monthly	<del></del>
Enteral / Pare		Dally		Calorie cou		No No	<u> </u>
		<del>-</del> ·	<u></u>				

-5

Dietitian Signature / Name / Date / Time:

11/1/2011:3

DATE AND TIME	DIETITIAN NOTES	SIGNATURE
N/1/24 11:30	A 59 years old gertlemen  came = clochest pain  was assessed to be cuell— nowwhed as evident by SGA.  K/C/O - No- co- mashidity  patient shifted to cathlob  for procedure (cop). bept on  NBM. patient recinel to  RL. NBM over recinel to	المحرك ر
11/1/24	an intotach soft solid diet  Educated ma prisent of  family on 1000 calarises, Low fat,  low salt diet on discharge  1000 ml fluid restricted  emphasized on Small  grequent neals & Down  Organic control.	\$28b
diet	hiet modifications à clarifications dons. Chart given on discharge	



59/Mule/MHI202481729 11/01/2024/IPH2024000093

Dr.G. GNANAVELU





## PATIENT TRANSFER FORM DIAGNOSTICS / PROCEDURES

Diagnosis:		NOTEMI/		1			NKOD.		
From (Area	a)	To (Area	1)	Date	Time	Reaso	Reason for Transfer / Name of Proce		
₽L-	er cath			11/1/24	11:15	5 (	Cary		
Method of Tra	Method of Transfer: ☐ On Bed ☐ On Wheelchair ☐ On Stretcher								
ASSESSMENT OF PATIENT: General condition of Patient: Conscious  Semi-conscious Un-conscious									
Language Ba	rrier: 🗌	Yes ☑ No ☐ If	Yes, spe	cify:					
Fall Risk Cate	gory: 🗆	Low Risk □ Me	dium Ris	k ☐ High F	≀isk				
Vital Signs (to l	be docur	mented at the tim	e of shift	ing):					
Temp (°F)	RR (	oreaths/min)	Puls	e (beats/mir	n)	SpO <sub>2</sub> (%)	BP (mmHg)	Pain	Score
97.2	a 2	Int	86	S/mt _		974.	117/80	0/2	ָ בֿס
Pain Scale used: ☐ PIPPS (28 weeks to ≤ 38 weeks) ☐ CRIES (38 weeks - 2 months) ☐ FLACC Scale (2 months - 7 years) ☐ Wong-Baker FACES Pain Rating Scale (7 years - 12 years) ☐ Numerical Rating Scale (>12 years) ☐ CPOT (ventilator / comatose)  Any pre-medication given: ☐ Any critical information: ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐									
	Sign	ature	Nar	ne			Emp. No.	Date	Time
Handover by	<b>y</b>	80		Sceet the	à		2002	11/1/24	1940
Handed over to	<u> </u>	<u></u>		Score the Sand	hoya	1. R	0004	11/1/24	11:20
After Procedure:  Procedure completed: ✓ Yes   Any critical information: ✓/  Vital Signs (to be documented at the time of shifting):									
Temp (°F)		oreaths/min)	Puls	e (beats/mir	n)	SpO <sub>2</sub> (%)	BP (mmHg)	Pain	Score
97.88	226	ช/ทเก	93.	beats/mi	n	95%	117/60	0/10	
Pain Scale used: ☐ PIPPS (28 weeks to ≤ 38 weeks) ☐ CRIES (38 weeks - 2 months) ☐ FLACC Scale (2 months - 7 years) ☐ Wong-Baker FACES Pain Rating Scale (7 years - 12 years) ☐ Wumerical Rating Scale (>12 years) ☐ CPOT (ventilator / comatose)									
	_ <del></del>	ature	Nan	11011			Emp. No.	Date	Time
Handover by	- /		8	and higa	· R		0004	11/24	12:20
Handed over to	o	TK\$ 1.50		Javan	ja.		DI 52 10	W1/24	12.20

The way to better health (A Unit of United Alliance Healthcare Pvt Ltd)

Mr.SAMRUDDIN S M

59/Malc/MHI202481729 11/01/2024/iPH2024000093

Dr.G. GNANAVELU





## **CONSENT FOR CORONARY ANGIOGRAM / CORONARY ANGIOPLASTY**

Dr CINAN AUTO. has explained that I have the following condition:

Fat (cholesterol) and calcium can build up in the arteries like rust in old pipes. It can stop the flow of blood to the heart. This can cause angina or a heart attack. The Coronary Angiography procedure is performed to show up the amount of disease in the coronary arteries, the blood vessels that supply the heart with bleed. After an injection of local anaesthetic, a fine tube (catheter) is put into the artery in the groin/hand. The tube is carefully passed into each coronary artery in turn. A series of video pictures are taken using xrays and an iodine containing contrast medium (x-ray dye). The contrast medium may be injected into the main pumping chamber of the heart (left ventricle). This helps us to find out whether you have any narrowing or blockage of your coronary arteries. The doctor can then tell you which treatment is best for you after carefully studying and discussing your pictures. This may be an operation such as a coronary by-pass or a procedure called an angioplasty (the arteries are widened using a small sausage shaped balloon). Sometimes, drugs alone may be a suitable option.

#### RISKS OF THIS PROCEDURE

The risk of coronary angiography depends on:

(i) The nature of coronary artery disease (ii) The pumping status of the heart (iii) Your age and general health These are some of the more serious risks that can happen, but are not the only risks:

Less than 1 in 10,000 (0.0001%)	(a) skin injury from radiation, causing, reddening of the skin
1 in 1000 people (0.001%)	<ul> <li>(b) A stroke. This can cause paralysis and long term disability</li> <li>(c) Heart attack.</li> <li>(d) A dangerous reaction to the x-ray contrast medium (dye). If this happens, you may have severe reactions such as asthma, shock and convulsions. Death in extremely rare cases about 1 in 2,50,000 to 4,00,000 injections.</li> <li>(e) Need for major surgery to the leg at the puncture site.</li> <li>(f) Need for emergency heart surgery or angioplasty.</li> <li>(g) A higher lifetime risk from x-ray exposure.</li> <li>(h) Death</li> </ul>
1 in 100 people (0.01%)	<ul> <li>(I) the heart may not beat in a proper rhythm which will need urgent treatment</li> <li>(j) Surgical repair of the groin puncture site. This may need a longer stay in hospital.</li> <li>(k) Minor reaction to contrast medium such as hives.</li> <li>(l) Loss/impairment of kidney function due to the contrast medium</li> </ul>
1 in 20 people (0.05%)	(m) Major bruising or swelling at the groin punture site
Most People	(n) Minor bruising

PATIENT CONSENT:

P acknowledge that Dr. Only No. 10. has explained my medical condition and the proposed procedure. I understand the risks of the procedure, the anaesthetic including the risks that are specific to me and the likely outcomes if complications occur. The Doctor has explained other relevant treatment options their risks and my right to refuse the treatment. He has explained my prognosis and the risks of not having the procedure. I have been explained that some reprocessed items might be used once its sterility and integrity is confirmed. I was able to ask questions and raise concerns with the doctor about my condition, the procedure and its risks, and my treatment options. My questions and converns have been discussed and answered to my satisfaction. I understand that in the unlikely event of complications, I may require a blood transfusion, an additional procedure or surgery. The doctor has explained to me that if immediate life-threatening events happen during the procedure, they will be treated accordingly. I understand that no guarantee has been made that the procedure will improve the condition. On the basis of the above statements,

#### I REQUEST TO HAVE THE PROCEDURE

	Signature	Name	Date	Time
Patient/Guardian with relationship	100 350n	HR. SAMRUDDIN.	11/1/24	LO- 38
witness	1(ED CO B)0	DARS. MUNHAU.	11/24	10.30
Doctor	\$ 93337 A	Dr-leAD THICLE	11/1/24	10:20
Interpreter			7	







Patient Details *(Affix Label here)* Name:

UHID: DOB: Sex:

## இருதய ஆன்ஜியோகிராம் பரிசோதனைக்கான ஒப்பம்

#### நீலை மற்றும் செயல்முறை

பழைய இரும்புக் குழாய்களில் துருபிடிப்பதைப் போல், தமனிகளில் கொழுப்பு மற்றும் கால்சியம் சேகும். இது ஆன்ஜினா அல்லது மாரடைப்பினை ஏற்படுத்துகிறது. இதயத்திற்கு ரத்தத்தினை வழங்கும் ரத்தக்குழாயான இதயச்சுவர் சிரை தமனிகளில் நோயின் அளவினை கண்டறிய கரோனரி ஆஞ்சியரேகிராஃபி செயல்முறை மேற்கொள்ளப்படும். ஒரு மோக்கல் அன்றத்படிக் முயக்க மருந்து) வழங்கப்பட்ட பின், ஒரு சிறிய குழாயானது (கதீட்டர்) கவட்டை/கையினுள்ள தமனியில் செனுத்தப்படும், இந்த குழாய் ஒவ்வொரு இதயச்சுவர் சிரை தமனிகளிலும் மாற்றி மாற்றி கவனமாக வரிசையாக செனுத்தப்படும். எக்ஸ்ரே மற்றும் பிற அயோடின் கொண்டுள்ளன காண்டுராஸ்ட் மீடியத்தினை (எண்ஸ்ரே டை) பயன்படுத்தி, பல வீடியோ படங்கள் வரிசையாக எடுக்கப்படும். இதயத்தின் மூக்கிய ஏற்றியிறைத்தல் அறையில் (இடதுபக்க இருதய கீழறை) இந்த கான்பராஸ்ட் மீடியம் உட்செனுத்தப்படலாம். இது இதயத்தின் அளவினை மதிப்பிடவும் மற்றும் அது எவ்வாறு பம்ப் செய்கிறது என்பதை மதிப்படவும் மேற்கொள்ளப்படும். இப்படங்கள் நமக்கு இதயச்சுவர்சிரை தமனிகள் குறித்த ஒரு படத்தினை வழங்கும். இது உங்களுக்கு ஏதேனும் அடைப்பு இருக்கிறதா என்பதை கண்டறிய உதவும். பின்னர் உங்கன் படங்களை கவனமாக பார்த்த பின் மருத்துவரால் உங்களுக்கு ஏற்ற சிகீட்சையை மேற்கொள்ள முடியும். இடையை படிக்கு கொண்டு தமனியை அகைப்படுத்துகல்) என்னும் ஒரு செயல்முறையாகவும் இருக்கலாம். சில நேரங்களில் மருந்துகள் மடிடுமே போதுமானதாக இருக்கலாம்.

#### கீச்செயல்முறையிலுள்ள கீடர்பாடுகள்

இதயச்சுவர் சிரை ஆன்ஜயோகீராஃபியிலுள்ள இடர்பாடுகள் பின்வருபவைகளையே சார்ந்திருக்கும்

(i) இதயச்சுவர் சிரை தமனி நோயின் தன்மை — (ii) இதயத்தின் ஏற்றியிறைத்தல் நிலை — (iii) இதயத்தின் வயது மற்றும் பொது ஆரோக்கியம் ஒற்பட வாய்ப்புள்ள சில தீவிர கேடர்பாடுகள் பின்வருமாறு. ஆனால் கிலைகள் மட்டுமே முழுமையான கிடர்பாடுகள் அல்ல

10,00-ல் ஒருவருக்கும் கீழ் (0.0001 சதவிக்தம்)	(a) கதிர்வீச்சின் காரணமாக ஏற்படும் தோல் பாதிப்பு, சருமம் சிவந்து போதல்
1000-ல் ஒருவருக்கு (0.001 சதவிகீதம்)	(b) வலிப்பு. இது பக்கவாதம் மற்றும் நீண்டநாள் ஊனத்தை ஏற்படுத்தலாம் (c) மாரடைப்பு (d) எக்ஸ்-ரே காண்ட்ராஸ்ட் மீடியத்தின் (டை) ஆபத்தான விளைவுகள் . இவை ஏற்பட்டால் உங்களுக்கு ஆஸ்துமா, அதீர்ச்சி மற்றும் வலிப்பு போன்றவைகள் ஏற்படலாம். 2,50,000 முதல் 4,00,000 ஊசிகளில் ஒன்று மரணத்தையும் வீளைவிக்கலாம். (e) குத்தப்பட்ட இடத்தில் பெரிய அறுவை சிகீச்சை மேற்கொள்ள வேண்டியது வரலாம். (f) அவசரகால இதய அறுவை சிகீச்சை அல்லது ஆன்ஐயோபினாஸ்டிக் தேவைப்படலாம். (g) எக்ஸ்ரே கதீர் பாதீப்பு காரணமாக அதிக வாழ்நாள் அச்சுறுத்தல் இடர்பாடு. (h) இறப்பு
100-ல் ஒருவருக்கு (0.01 சதவிகீதம்)	<ul> <li>(I) இதயம் சரியான முறையில் துடிக்காமல் இருக்கலாம். அதற்கு அவசரமாக சிகீச்சை தேவைப்படு:</li> <li>(j) குத்தப்பட்ட கவட்டை பகுதியில் அறுவை சிகீச்சை சரிபாடு. இதனால் மருத்துவமனையில் நீண்ட நாட்கள் தங்கீயிருக்க வேண்டியது வரலாம்</li> <li>(k) தோல் அரிப்பு போன்ற சிறு விளைவுகள்</li> <li>(l) கான்ட்ராஸ்ட் மீடியம் காரணமாக சிறுநீரகம் செயல்படாமை அல்லது அதன் வலுகுறைதல்</li> </ul>
20-ல் ஒருவருக்கு (0.01 சதவிலிதம்)	(m) குத்தப்பட்ட இடத்தீல் பெரிய அளவிலான சிராய்ப்பு அல்லது வீக்கம்
பெரும்பாலான மக்களுக்கு	(n) சிறிய அளவிலான சிராய்ப்பு

#### நோயாளி ஒப்புதல்

#### செயல்முறையை எனக்கு மேற்கொள்ளுமாறு கேட்டுக்கொள்கிறேன்

	கையெழுத்து	பெயர்	தேதி	நேரம்
நோயாளி (பாதுகாவலர்) உறவுமுுசுற		•		
சாட் <b>சி</b> ்				
மருத்துவர்	4 '			
மொழிபெயர்ப்பாளர்			•	







## TRANSRADIAL CORONARY ANGIOGRAM REPORT

Patient Name:	Mr. SAMRUDDIN.S.N	ID:	MHI202481729	
Age/Gender :	59 M	IPH:	IPH2024000093	
Cath No. :	3587		DOP:	11.01.2024
Done by	Assisted by	Technician	Phy	sician assistant
Dr.Gnanavelu/ Dr.Karthik	Ms. Bhavatharini	Mr. Ram Ms		Ms. Shalini

DIAGNOSIS: CAD- EVOLVED AWMI; COPD; MODERATE LV DYSFUNCTION

Access: Left radial artery

Total exposure time: 229.3"

Hardware used: 5F sheath, 5F TIG

Total DAP: 26.71 Gy.cm<sup>2</sup>

Contrast used: CONTRAPAQUE 40 ml

Total RAK: 123,44 mGy

Medications given: Inj NTG 200 mcg & Inj Heparin 2500 IU IA

Hemodynamic data: Aortic pressure: 124/80(95) mmHg; HR 91 bpm; SpO2 96%

Selective Coronary angiogram done in multiple angulated views:

ARTERY	FINDINGS
LEFT MAIN	Normal. Bifurcates into LAD & LCx.
LAD	Type 3 vessel. Ostial LAD has near total occlusion with TIMI I distal flow.  Major diagonal is a large vessel, has diffuse ostioproximal disease with TIMI  Il distal flow. Mid and Distal LAD visualized by heterocollaterals.
LCx	Nondominant. Proximal and distal LCX have luminal irregularities. Gives 2 major OMs which have luminal irregularities.
RCA	Dominant. Proximal RCA has luminal irregularities. Mid and Distal RCA are normal. PDA and PLv have luminal irregularities.
LIMA	Normal.

FINDINGS: RIGHT DOMINANT SYSTEM; OSTIAL LAD NEAR TOTAL OCCLUSION

ADVICE: IVUS GUIDED PTCA X LAD (POBA X DIAGONAL) vs CABG

Dr. G.GNANAVELU, MD, DM.

Dr. G. Gnanavelu MD, DM (cardio), FACC Chief Cardiologist Reg. No: 39469

#9, 1st Main Road, United India Colony, Kodambakkam, Chennai - 600024. Tel : 044 - 4310 8959

**★** @MedwayHospitals

(a) @medwayhospitals

In @medway-hospitals

@medwayhospitals

94557 94557 1800 572 3003

Medway Group of Hospitals

Medway Centre of Excellence (Chennai)



59/Malc/MHI202481729

11/01/2024/IPH2024000093

Dr.G. GNANAVELU



#### MHI/NUR/2022/048

DATE & TIME		Observation / Action			Signature with Emp.No
11/1/24	⇒ Patien!	5 got adr	nicefon	ND ND	
		stocke for			
	CAGI. ta	atient was	chain	$\mathcal{D}$	- Paige
	dumme cell	, Stable.	Vitale		0.0
	noemal -	Sarge.			·
	=> NPO	maintained	@ 9,	00	
	=> propo	wation de	me		
		-faken		·	
	→ Tv	cline celle	cortad		70
11:15	-> Patie	It. Shifte		<u>) ;                                     </u>	10 (34
	cathlab	@ 11:15			
1-1-2-4-		CATHLAB REPO	RZS:	<u></u>	
11/1/	e patient r	cived from RL	to cathe	lab	<i>-</i>
11.20	Phis Conscious and good oriented Juline polant  Phis Conscious and good oriented Juline polant  Phis Continuously Cardiac Monitoring HE 9  Phis Continuously Cardiac Monitoring HE 9				¥3004
			Moritona	9 4/29	26pm.
(1, '	Bp-120/85(47)	8 p.o 2 95%			
11/4	Bp-120/85047) Bp-120/85047) BSkrille dro	pping done.	JUF N. 30	11/map	، سر
	& CAU procedi	we start th	rough Lef	1 Radia	- P
11. 57	artery approa	ch under Glocal	arestles "	7,00	6004
.,,,	Dung pro	ch under Grocal. Ceclure Gr. NTU 500 unib Agui	200m(U)	nd	
12.	G. Hepwin dr	bround JAgiu	en- B/0-Di	r. E.S Sa	
2:10	D IVF. NB	500 unis Agin 30n1 /hr reflow 08mal : HR-91 got over Pf	s pt B.	Stable	
2.	vitals alera	Smal. HR-91	1/21 ada	0-117160	Spor 95%
12'		yot over. Pt	U. hemod	grille	
	Stable - Signature	Name	Emp . No.	Date	Time
Document		Ivalite	cmp. No.	Date .	inile
endorsed by	· · · · · ·	Sandhiya-R	0004	11/1/20	4 12:46



institute where

DATE &		Observation / Action			Signature with Emp.No
11/25	10H Rads	al astery shea	th romo	reel	
12:15	mel 19ght	prosence pando	safe applic	el. no	6)
, ' , 1	11071 (11610	TUTTO TO THE CONTRACT OF THE PARTY OF THE PA		· I)	2004
	& py Shiff	led to RLWith	UI doc	umont	
12:20	patient how	ding over to RE	W. Lawmyn	,	2,04
12.25		t secetived for		Rlab	<del></del>
<b> </b>	Battent	was shedr	wdynam	cally	TOLCO
, <u>, , , , , , , , , , , , , , , , , , </u>	Stable.	enoi done	, V fenc	lungs"	<del>-</del>
		an: pFCA-LI		0	··
<b> </b>	-> Had	fulce no	nause	a	
	Sensatio	y .	<u> </u>		108.
10.0	=> Solf => Patio	rolded.  Int chad i	0 0	0	
13:30		tiet.	sunch_		ouse.
18:10	hoemal > Pot	ient for a	Sidragard		
10.10		Semovos		10 d	
	fand xo	moved no	anu <u>~</u>	70	
	M&ML.	Vitals Stal	le THP	780	
		Committee =		/ 1	
	99	scharge adv	orco medi	ration	1- Kl
	explained	to batient	- she	2	0128.
	family	·			
 	'				
			1 - ·		
Document	Signature	Name	Emp . No.	Date	Time
endorsed by	Jay ooo	JAYADEVI-J	0002	rililer	18,30





## SAFE PROCEDURE CHECKLIST Adapted from WHO Safe Surgery Checklist



Mr.SAMRUDDIN S M

59/Malc/MHI202481729 11/01/2024/(PH202)

701/2024/IPH2024000093
Dr.G. GNANAVELU
<u> </u>

	•				. 1 /2	59/Malc/MHI20	2481729		
Name of the Procedure :	CAUI	Location :	CATHLAB-II	Date & Time :	01/2	11/01/2024/11	12024000093		
		4			e-1	1.15 Dr.G. GNANAVE	Lir		
Does the Procedure involve	Procedural Sedation : [_	] Yes [⊾Nb					iiiiiiiiiiiiiiiiiiiiiiiiiiiiiiiiiiiiii		
SIGN IN 1/ 30 Before Induction of Procedural Se	- d-41	TIME OUT // 4/	O Sedation and before procedure		SIGN OUT /	2, bo or indicates that the Procedure			
(Anaesthetist / Qualified Physicla		Atter production	(Anaesthetist or Qualified Physic	ian administering Procedu	I		13 completed		
Sedation + Nurse + Technician + Do	ctor performing the procedure)		<u> </u>	performing the Proce					
Patient Confirmation		All team members in	troduce themselves by Name and F	Role	To be done for procedures	or each procedure in case of n	· ,		
Identity by two identifiers	1) As	Identity by two identif	īers	□ Yes		Procedure done written down  (AUT)	Ľ <b>∃</b> Ýøs		
Procedure	□Xfes	Procedures	CA01	/ PYes		te of all specimens / investigat	ons □Yes⊡KVA		
Side	□Rł 🗹 Ll □NA	Side Loft Rad	tent goteny approals	PRI CELE/CINA	Confirms labe	eling and sent to lab	. /		
	1	Expected Blood loss					1		
Consent	<b>M</b> Yes ✓	Position Su	ip9 ne	LyY∕es	Any recovery		ĭ Yes ☐ None		
Known Allergy	□Yes □YN•	Consent	Taken	. □ <b>X</b> Yes	If Yes, Pis. s	pecify:			
!	If yes, plaese specify	Required equipment	and implants available	□ Yes □ NA					
Diff. It is a second of	Ma DVa andament	Essential Imaging dis	polavod	Miles □NA	obse	nursion.			
Difficult airway / aspiration risk / dentures	☑No ☐ Yes, equipment and assistance available		s within last 60 minutes	☐ Yes ☐ MA	-		ĺ		
Possibility of hypothermia	✓ No ☐ Yes, warmer in place	Name of the Antibioti			Any Equipme	ent / instrument problem that n	eeds to be		
Possibility of hypothermia	TINO T tes, waither in place		polism Prophylaxis Provided	☐ Yes ☑ NA	addressed:	•	□Yes □WAne		
All concerned anesthesia equipment a	and madication chack complete			□ Yes Ly Ny	if Yes, Pis. s	pecify:	/		
		Anticipated duration		+	4				
<del></del>	s pls. specify	Anticipated blood los		□X¶s □NA	4	_			
Pre OP medication taken	□Yes ☑⁄/)o	Adequate fluids and		□ Yes □ NA	ļ <u> </u>				
Bouled aguine at fac	□ Yes □ NA	For procedural sedat	critical or unexpected steps	□Xŧs	Corrective act	tion :			
Required equipment for procedure available	∐ Yes/ ∐ NA	Any patient specific of		☐ Yes ☐ Kine	-	/			
procedure available		Intra procedure glyce	ernic control	☐ Yes ☐ ÑĀ	_	/			
		Any concerns about	sterility	☐ Yes ☐/Ndne	1				
Anaesthetist / Doctor giving	Doctor performing to	ne & Nur	se: RN·Bara	Technician : 9/1 R	an	Others Please Specify :			
Procedural Sedation	Procedure :	4378	0 # 6		0001				
		125/1//	1124	1-110	), [				
Date :	Date: // /0//		e: 11/01/24	Date: 11/01/2	7	Date :			
Time :	Time: 12:10	Tim	e: /2!10	Time: /2,10		Time:			







Every heart beat counts

The way to better health (A Unit of United Alliance Healthcare Pvt Ltd)

#### **Procedure Monitoring Sheet (Cath Lab)**

Mr.SAMRUDDIN S M

Patient Nam 59/Malc/MHI202481729

11/01/2024/IPH2024000093

UHID / IP:

Dr.G. GNANAVELU

Consultant:

Ward Unit:

Als- Nyram Diagnosis:

## Pre Procedure Checklist (Please tick appropriately – To be filled by the Ward Nurse) **PARAMETERS** YES NO NA Vital signs: BP:117/80 Temp: 15. .... Pulse: 86... RR: 24/48P02: 49) Urine voided **Bowel preparation** Pre-procedure medication administered Procedure site marked Skin preparation done 9,00 **NPO** SON Loose Tooth removed Contact lenses / Eye glasses removed Prosthesis present Jewellery/Nail polish removed Checked for Allergies (Drug / food) IV line/In-situ Consent taken Investigation reports / Documents received Signature of Nurse: Date & Time : 1 Intra - Procedural Record (To be filled by the Cath Lab Nurse) Time HR / min RR / min BP mmHg SpO<sub>2</sub>% Medication / Remarks Sign. of Nurse 45% 95 % OVEY

## Post Procedure Follow Up Data (to be filled by the doctor) Route: Left Radial astery approach Complication : ~// BP: 106/10 mmHg, HR: <u>876pm</u>, RR: <u>20</u>, SpO2: <u>967</u> Distal Pulse: Yelt , Puncture Site: 100000109, 10 Lan toma Advise: Shift To: Ward / ICU ♦ Diet \_ (5 Inform Duty Medical Officer SOS a) If patient complains of any Discomfort b) If dressing is Loose or Socked with Blood c) If limbs\_are Cold / Absent Pulse Remove The banclage dressing on 12/1/24 at 11:00 AM /PM after informing to the consultant. Special instruction if any: Ni/ Name & Signature of Consultant POST PROCEDURE OBSERVATION SpO2% Date & Time HRIRR **Extremity Status** Remarks Sign. of Nurse Site Evaluation 11/1/24 Radial Goog No oras g 117/70 R-2004 96% 1 . . . . . . . Nurses Notes: CAOT procedure got over. Ptil Stable. Left Radial astery shorth removed and right pressure bandage applied. no orging no heretoma. Critical

Patient shift to: Recovery Room Patient Room CCU Other RL

Name & Signature of the Nurse: Roothinguisk Date & Time: 11/1/24

Smdhiguisk Date & Time: 11/1/24





Score Interpretation: Minimal Risk: 23 - 19; At Risk / Mild Risk: 18 - 15; Moderate Risk: 14 - 13; High Risk: 12 - 10; Severe Risk: 9 - 6

Mr.SAMRUDDIN S M 59/Male/MHI202481729 11/01/2024/IPH2024000093

I Dr.G. GNANAVELU





Date: BRADEN SCALE FOR PREDICTING PRESSURE INJURY RISK E Time: 4. No Impairment SENSORY 1. Completely Limited 2. Very Limited 3. Slightly Limited PERCEPTION -Presponds to verbal Unresponsive (does not moan, flinch,or Responds only to painful stimuli. Cannot Responds to verbal commands, but grasp) to painful stimuli, due to diminished communicate discomfort except by cannot always communicate discomfort commands. Has no sensory ability to respond meaning-fully to level of consciousness or sedation OR moaning or restlessness OR has a or the need to be turned OR had some deficit which would limit ability to feel or voice pain or pressure-related limited ability to feel pain over most of body sensory impairment which limits the ability sensory impairment which limits ability to discomfort to feel pain or discomfort over 1/2 of body feel pain or discomfort in 1 or 2 extremities discomfort 4. Rarely Moist 1. Constantly Moist 3. Occasionally Moist 2. Very Moist MOISTURE Skin is occasionally moist, requiring an Skin is usually dry, linen only Skin is kept moist almost constantly by Skin is often, but not always moist. Linen degree to which requires changing at routine extra linen change approximately once a perspiration, urine etc. Dampness is must be changed at least once a shift skin is exposed detected every time patient is moved or intervals to moisture turned 1. Bedfast 2. Chairfast 3. Walks Occasionally 4. Walks Frequently Confined to bed **ACTIVITY** Walks occasionally during day, but for very Walks outside room at least Ability to walk severely limited or non3 degree of existent. Cannot bear own weight and / or short distances, with or without twice a day and inside room physical activity must be assisted into chair or wheelchair. assistance. Spends majority of each shift at least once every two hours in bed or chair. during waking hours 4. No Limitation 3. Slight Limited 1. Completely Immobile 2. Very Limited MOBILITY Makes major and frequent Does not make even slight changes in body Makes occasional slight changes in body Makes frequent through slight changes in ability to change changes in position without or extremity position without assistance or extremity position but unable to make body or extremity position independently and control body assistance frequent or significant changes position independently 3. Adequate 1. Very Poor 2. Probably Inadequate 4. Excellent Eats over half of most meals. Eats a total of Never eats a complete meal. Rarely eats Rarely eats a complete meal and generally Eats most of every meal. more than any food offered. Eats 2 servings eats only about 2 of any food offered. 4 servings of protein (meat, diary Never refuses a meal. NUTRITION or less of protein(meat or dairy products) per Protein intake includes only 3 servings of products) per day. Occasionally will refuse Usually eats a total of 4 or usual food a meal, but will usually take a supplement day. Takes fluids poorly. Does not take a meat or diary products per day. more servings of meat and intake pattern liquid dietary supplement OR Is NPO and / or Occasionally will take a dietary when offered OR is on a tube feeding or diary products. Occasionally maintained on clear liquids or IV's for more TPN regimen which probably meets most eats between meals. Does supplement of nutritional needs than 5 days not require supplementation 1. Problem 2. Potential Problem 3. No Apparent Problem Moves in bed and in chair independently and has sufficient muscle Requires moderate to maximum assistance Moves feebly or requires minimum in moving. Complete lifting without sliding assistance. During a move skin probably strength to lift up completely during move. Maintains good position in bed **FRICTION** against sheets is impossible. Frequently slides to some extent against sheets. orchair & SHEAR slides down in bed or chair, requiring chair, restraints or other devices. **TOTAL SCORE** frequent re-positioning with maximum Maintains relatively good position in chair assistance. Spasticity, contractures or or bed most of the time but occasionally Initial & Emp. No. agitation leads to almost constant friction slides down of Staff Nurse: Initial & Emp. No.

of Sr. Staff Nurse:





59/Male/MHI202481729 11/01/2024/IPH2024000093

Dr.G. GNANAVELU

## Dr.G. GNANAVELS


MHI/NUR/2022/052



Every heart beat counts

## **PAIN RE-ASSESSMENT & MONITORING CHART**

Date & Time	Pain Score	Pain Character (dull, achy, sharp, stabbing, shooting, burning, referred / radiant pain)	Duration	Location / Site	Interventions	Staff Initial & Emp. No.	Senior Staff Initial & Emp. No.
111124	9/60	No pour	1		·	AL 70158	Jan John
11.15	olw	No pain				70158	Jaros
12:85	- 0/u	6 /	P+ (	shifted s	to eath (a) 11.35.	Pisa	Jan (88)
13.25	WV	No pain	1	_		PLSQ	Joe Jose
1425	Vío	No pain		(		92.58	Jan 600
15.25	0/10	No pain				0158	Jon Lago
16:25	Oùo	•				POLCA	Jul 280
[ <del>]</del>	- Wo	No pain				2158	Jul 600
		Pali	ent	got a	Rechauged 918,10.		







59/Malc/MHI202481729 11/01/2024/IPH2024000093

Dr.G. GNANAVELU





## **DVT RISK ASSESSMENT**

Ass	ign a score of 1 if (YES) in parameter nos. 1 to 9,	and ass	ign a sc	ore of -2	if (YES)	in parai	neter no	. 10
	Date	11/1/20						
		10115				<u> </u>		
S. No.	PARAMETERS		_					
1	Active cancer (on-going treatment or diagnosed within 6 months or palliative care)	0			_			
2	Bedridden recently >3 days or major surgery within four weeks	0						
3	Calf swelling >3 cm compared with asymptomatic side, measured at 10 cm below tibial tubercle (Assess for both legs)	0						
4	Collateral (nonvaricose) superficial veins present (Assess for both legs)	0			l			
5	Entire leg swollen (Assess for both legs)	0_						
6	Localized tenderness along the deep venous system (Assess for both legs)	0						
7	Pitting edema, greater in the symptomatic leg (Assess for both legs)	0						
8	Paralysis, paresis, or recent plaster immobilization of the lower extremity (Assess for both legs)	0						
9	Previously documented DVT (Assess for both legs)	0_						
10	Alternative diagnosis to DVT as likely or more likely (Assess for both legs) / Co-morbidity like ESLD / Renal disease, Renal failure, CCF Cellulitis (commonly mistaken as DVT), Dependent (stasis) oedema, Lymphatic obstruction. Septic arthritis, Cirrhosis, Nephrotic syndrome, Calf muscle tear or strain, Haematoma (collection of blood) in the muscle, Sprain or rupture of a leg tendon, Fracture.	0						
	FINAL SCORE	0						
Low F	lisk: -2 to 0   Moderate Risk: 1 to 2   High Risk: 3 to 8	was						
	DVT prophylaxis started	☐ Yes ☑ No	□ Yes □ No	□ Yes □ No	☐ Yes ☐ No	☐ Yes ☐ No	□ Yes □ No	☐ Yes ☐ No
	Signature & Emp. No. of RN	10150						
	Signature & Emp. No. of Sr. RN	< 12/2 Pr	,					



## Medway Hospitals

The way to better health
(A Unit of United Alliance Healthcare Pvt Ltd)



Mr.SAMRUDDIN S M 59/Male/MHI202481729 11/01/2024/PH2024000093

Dr.G. GNANAVELU





## MODIFIED MORSE FALL RISK ASSESSMENT CHART

	· ;	i -	··· , ·						. : 
Date	11/1/20	11/1/2						, ,	
Time	10.15	13.25				values bu	. "-	6 4 396 13 .	1'
No	0	(3)	ò	, Q	້0	0	Ó	Ö	Ó
Yes	25	. 25	·25	25	25	25	25	25	25
. No	.(0)	(0)	0	0 _	0	0.	_ 0 ' :	0	1 0
Yes	15	15	15	15 <sup>-</sup>	15	15	<del>7,</del> 15	15	15
No	0	0	0	0	0	0	0	.0 .	0
Yes	(20)	(20)	20	20	20	20	20	20	20
		) (	1					£ 1jr - 1	
	(0)	(0)	0	0	0	0 .	0	<u> </u>	. 0
	15-	15	15	15	-15	15	15	15	15
	30	30	30	30 📆	30	30	30	30	30
<u> </u>		3	1 :15	.a 1565	, ,	۱ .		•	7
- '	(O)	(0)	<b>O</b> ,	, <b>0</b> ,	0	- <b>0</b> -	0	0	0
- :	10	10	10			, vi10 1 in	P. 1.0	10,	10
# # # # # # # # # # # # # # # # # # #	20	20 :	<b>20</b> %			320 <sup>m</sup>	# <b>20</b> <sup>(4)</sup> -	) <b>20</b> :	20
j i				i zi	ا اا برر	- - 10		الأوا	
	(0)	_(0)	0	0	. 0	0	0	.0	· O
	- 15	15	15	-15	<b>15</b>	Жа <u>а</u> ты ыл. - <b>15</b> -	-15	15	-15
									•
No	(0)	(6)	0	0	0	0	; i, <b>0</b> ; ;		٠,0
Yes	15	15	15	~. <b>15</b> .	15	15	15	15	-15
	:		·	, ;- · · · · · · · · · · · · · · · · · ·	•	~		1 - ' 1	
,	20	as.	وداره يې ؤ	T T 14 1 21	'; '			, 1	
- ' .	,				1551	grijerini i	3	e ije i j	
t	,	<u>'</u>	4 .			100 40	100	• 1	
	SL AV	Ha	<u>ार्</u> ग औ	- ;		97 C. 2 pts	20° £	· 10	11 ;
	10.72	~	:				1		
	Time No Yes No Yes No Yes No Yes	Time No (0) Yes 25 No (0) Yes 15 No (0) Yes (20)  15 30 10 20 15 No (0) Yes 15  No (0) Yes 15	Time No 0 0 0 Yes 25 25 No 0 0 0 Yes 20 20 20 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	Time No 0 0 0 0 0 Yes 25 25 25 No 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	Time No 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	Time No	Time No	Time No (0) (0) (0) (0) (0) (0) (0) (0) (0) (0)	Time No 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0

INTERVENTIONS  Titok aper the Risk Score  Time  10:15  Low Risk Interventions (0 - 24)  Familiarize the patient with the immediate surroundings  Remind the patient to use call believing exting out of ted.  Keep the two sider alia in the raised position at all times for algaliatins regardless of age.  Keep the two sider alia in the raised position at all times for patients easy reach.  Remove excess equipment or furniture to make a clear path.  Keep the patients bed in the low position at all times except outing procedure.  Remove excess equipment or furniture to make a clear path.  Remove excess equipment or furniture to make a clear path.  Remove excess equipment or furniture to make a clear path.  Remove excess equipment or furniture to make a clear path.  Remove excess equipment or furniture to make a clear path.  Remove excess equipment or furniture to make a clear path.  Remove excess equipment or furniture to make a clear path.  Remove excess equipment or furniture to make a clear path.  Remove excess equipment or furniture to make a clear path.  Remove excess equipment or furniture to make a clear path.  Remove excess equipment or furniture to make a clear path.  Remove excess equipment or furniture to make a clear path.  Remove excess equipment or furniture to make a clear path.  Remove excess equipment or furniture to make a clear path.  Remove excess equipment or furniture to make a clear path.  Remove excess equipment eraised point and and the excess of the medications that effects level or or social path.  Remove excess equipment eraised point and and the excess of the medications that effects level or consciousness, gait and elimination when planning patients care  Do not leave patients unattended in diagnostic or treatment areas.  Remove experiment excess the path of the pa		Date	1,/25	1118			}			-	
Low Risk Interventions (0-20)  Familiarize the patient with the immediale surroundings  Reamind the patient to use call bell before getting out of bed  Keep the two side rails in the raised position at all times for all patients regardiess of age  Keep the call bell, betakide table, water, glasses within the patients easy reach  Remove excess equipment or furniture to make a clear path  Keep the patients ted in the low position at all times except during procedure  Reach fall prevention techniques, such as sitting up for a moment before fising from the bed  Bed whosels should be locked  Ensure that floor of the bathnoomis dry and not slippery  Review medications for potential side effects that can promote falls  Des safety belts during movement in wheelchair  The patients are not ambulated by themselves. They are to be ambulated orly with assistance  Recitamistification and in the patients are instituted for heavy or delimited patients in a bed or wheel chair or on a toller seal.  Lose restricts and bed monitions as ordered by the doctor.  Allow the patient to embulate only with assistance  Consider peak effects of the medications that effects level or consciousness, grait and elimination when planning patients care  On not cleave patients unattended in diagnostic or treatment areas  Accompany the patient to use grab bars near the tollet, bathtub, and shower  Make sure the family and other visitors understand the restrictions mentioned above  [Igh-risk/Interpentions (25 or above)]  Apply all the low and medium risk interventions  The rest all risk up in the bod whoch chart and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Frovide a commode at bedside (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety helds	INTERVENTIONS	- Date	Hick	Mu		_					
Familiarize the patient with the immediate surroundings Hernind the patient to use call bell before getting out of bad. Keep the two side rails in the raised position at all times for all patients regardless of age. Keep the call bell, bedside table, water, plasses within the patient's easy reach Path search seasy reach Path se	Tick as per the Risk Score	Time	10:15	12.2	5						
Familiarize the patient with the immediate surroundings Hernind the patient to use call bell before getting out of bad. Keep the two side rails in the raised position at all times for all patients regardless of age. Keep the call bell, bedside table, water, plasses within the patient's easy reach Path search seasy reach Path se	Low Risk Interventions (0 - 24)				•				_		
Remind the patient to use call bell before getting out of bad.  Keep the two side ralls in the raised position at all times for all patients regardless of age.  Keep the call bell, bedside table, water, plasses within the patients easy reach.  Remove excess equipment or furniture to make a clear path.  Keep the patients bed in the low position at all times except during procedure.  Teach fell-prevention techniques, such as sitting up for a moment before rising from the bed.  Bed wheels should be locked.  Encourage family participation in the patient's care promote falls.  Les safety bets during movement in wheelchair.  The patients are not anulutated by themselves. They are to be ambutated or ly with assistance.  Medium iskinterventions (25-44).  Apply all the low risk interventions.  Tie yellow fall risk tag in the bed and Wheel chair / Strocther.  Make sure that proper transfer precautions are instituted for heavy or debilitated patients in a bed or wheel chair or on a tollet seat.  Use restraints and bed monitors as ordered by the doctor.  Allow the patient to ambutate only with assistance.  Do not leave patients unattended in diagnostic or treatment areas.  Do not leave patients unattended in diagnostic or treatment areas.  Accompany the patient to ambutate only with assistance.  Do not leave patients unattended in diagnostic or treatment areas.  Accompany the patient while going to bathroom.  Advice the patient to segrab bars near the toilet, bathtub, and shower.  Make sure the family and other visitors understand the restrictions mentioned above.  If provide a commonded the diagnostic or freatment areas.  Apply all the low and medium risk interventions.  The restrictions mentioned above.  If proprietal, consider using protection devices: safety better.  Britant and the patient search of the properties.  Britant and the patient search of the properties.  Encourage tamily members or other visitors to stay with them.		ings	اس ا	〜						•	
Keep the call belt, bedside table, water, glasses within the patient's easy reach Reap the call belt, bedside table, water, glasses within the patient's easy reach Remove excess equipment or furniture to make a clear path Keep the patient's bed in the low position at all times except during procedure Teach felt-prevention techniques, such as sitting up for a moment before rising from the bed Bed wheels should be locked Encourage family participation in the patient's care Ensure that floor of the bathroom is dry and not ellippery Review medications for potential side effects that can promote talls Uses selety belts during movement in wheelchair The patients are not ambutated by themselves. They are to be ambutated only with assistance Rediturn task interventions? Tile yellow fall risk tag in the bed and Wheel chair / Stretcher Make sure that proper transfer precautions are instituted for heavy or debilitated patients in a bed or wheel chair or on a folial seal. Use restraints and bed monitors as ordered by the doctor Allow the patient to ambutate only with assistance  On not leave patients unattended in diagnostic or treatment areas Accompany the patient while going to bathroom Advice the patient to use grab bars near the toilet, bathub, and shower Make sure that family and other visitors understand the restrictions menomed as bedsite interventions Tile relation to use grab bars near the toilet, bathub, and shower  Make sure the family and other visitors understand the restrictions menomed as bedsite interventions Tile relation to use grab bars near the toilet, bathub, and shower  Interventions (45 or above)  Apply all the low and medium tilsk interventions Tile relation to use grab bars near the toilet, bathub, and shower  Interventions (45 or above)  Answer these patients call bella sa quickly as possible  Provide a commode at bedsite (frappropriate)  Utrinative these patients call bella sa quickly as possible  Encourage family members or other visitors to stay with then			•			_		i			
Keep the call bell, bedside table, water, glasses within the patient's easy reach Remove excess equipment or furniture to make a clear path Keep the patient's bed in the low position at all times except during procedure  Teach fail-prevention techniques, such as sitting up for a moment before rising from the bed Bed wheels should be locked  Encourage family participation in the patient's care Ensure that floor of the bathroomis dry and not slippery Review medications for potential side effects that can promote falls Use safety belts during movement in wheelchair The patients are not ambulated by themselves. They are to be ambulated only with assistance  Retirm risk interventions (256-44) Apply all the low risk interventions  Tie yellow fall risk tag in the bed and Wheel chair / Stretcher Make sure that proper transfer precautions are instituted for heavy or debilitated patients in a bed or wheel chair or an tollet seat Use restraints and bed monitors as ordered by the doctor Allow the patient to ambulate only with assistance  Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patients care  Do not leave patients unattended in diagnostic or treatment ransa Accompany the patient while going to bathroom Advice the patient to use grab bars near the toilet, bathub, and shower  Make sure the family and other visitors understand the restrictions mentioned above we are strictions. The restrictions mentioned above we may be a surface of the patients care.  John Patient of the patients in a room olose to the nurses' station.  Answert these patients call bells as quickly as possible Provide a commode at bedside (flappropriate)  Linakbedpan should be within easy reach (flappropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts									<del>-</del>		
patient's easy reach Hemove excess equipment or furniture to make a clear path Keep the patient's bed in the low position at all times except during procedure Teach fall-prevention techniques, such as sitting up for a moment before rising from the bed Bed wheels should be locked Encourage family participation in the patient's care Ensure that floor of the bathroom is dry and not slippery Review medications for potential side effects that can promote falls Use safety betts during movement in wheelchair The patients are not ambulated by themselves. They are to be ambulated only with assistance Meditim risk interventions They altow fall inskitag in the bed and Wheel chair / Stretcher Make sure that proper transfer precautions are instituted for heavy or debilitated patients in a bed or wheel chair or on a tollet seet. Use restraints and bed monitors as ordered by the doctor Allow the patient to ambulate only with assistance Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patients care Do not leave patients unattended in diagnostic or treatment areas Accompany the patient while going to bathroom Advice the patient to use grab bars near the toilet, bathrub, and shower Make sure the family and other visitors understand the restrictions mentioned above [Igh-Irsk interventions (45 or above) Igh-Irsk interventions (45 or above) Igh-Irsk interventions (45 or above) Answer these patients call belis as quickly as possible Answer these patients call belis as quickly as possible Provide a commode at bedside (if appropriate) Ulrinal/bedpan should be within easy reach (if appropriate) Irr propriate, consider using protection devices: safety belts  Signature & Emp. No. of RN	all patients regardless of age					1					
Remove excess equipment or furniture to make a clear path  Keep the patient's bed in the low position at all times except during procedure  Teach fall-prevention techniques, such as sitting up for a moment before rising from the bed  Bed wheels should be locked  Encourage family participation in the patient's care  Ensure that floor of the bathroom is dry and not slippery  Review medications for potential side effects that can promote falls  Use safety betts during movement in wheelchair  The patients are not ambulated by themselves. They are to be ambulated only with assistance  **Medicum risk interventions**  Tre yellow fall risk flag in the bed and Wheel chair / Stretcher  Tre yellow fall risk flag in the bed and Wheel chair / Stretcher  Wake sure that proper transfer precautions are instituted for heavy or debilitated patients in a bed or wheel chair or on a toilet seat  Use restraints and bed monitors as ordered by the doctor  Allow the patient to ambulate only with assistance  Consider pask effects of the medications that effects level of consider pask effects of the medications that effects level of consider pask effects of the medications that effects level of consider pask effects of the medications that effects level of consider pask effects of the medications that effects level of consclousness, gait and elimination when planning patients care  Do not leave patients unattended in diagnostic or treatment areas  Accompany the patient while going to bathroom  Advice the patient to use grab bars near the toilet, bathtub, and shower  Make sure the family and other visitors understand the restrictions mentioned above  Infinity	Keep the call bell, bedside table, water, glasses w	ithin the	1	/					-		
path Keep the patient's bed in the low position at all times except during procedure Teach fall-prevention techniques, such as sitting up for a momentate fore sing from the bod Bed wheals should be locked Encourage tamily participation in the patient's care Ensure that floor of the bathroom is dry and not slippery Review medications for potential side effects that can promote falls Use safety belts during movement in wheelchair The patients are not ambulated by themselves. They are to be ambulated only with assistance (Medium risk interventions (25-41) Apply all the lond wisk interventions Tie yellow fall risk tag in the bed and Wheel cheir / Stretcher Make sure that proper transfer precautions are instituted for heavy or debilitated patients in a bed or wheel chair or on a tollet seat Use restraints and bed monitors as ordered by the doctor Allow the patient to ambulate only with assistance Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patients care Do not leave patients unattended in diagnostic or treatment areas Accompany the patient while going to bathroom Advice the patient to use grab bars near the toilet, bathrub, and shower Make sure the family and other visitors understand the restrictions mentioned above Inglies/Sinterventions (45 or 850 vc) Apply all the low and medium risk interventions Tie red fall risk tag in the bed, wheel chair and stretcher Locate the high-risk patients in a room close to the nurses' station Answer these patients call bells as quickly as possible Provide a commode at bedside (if appropriate) Urinat/bedpan should be within easy reach (if appropriate) Fineourage family members or other visitors to stay with them If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN	patient's easy reach									ļ	
Keep the patient's bed in the low position at all times except during procedure  Tacah fall-prevention techniques, such as sitting up for a moment before rising from the bed Bed Wheels should be locked  Encourage family participation in the patient's care  Ensure that floor of the bathroom is dry and not slippery  Review medications for potential side effects that can promote falls  Use safety belts during movement in wheelchair  The patients are not ambulated by themselves. They are to be ambulated only with assistance  Weefflum risk finiterventions (25-44)  Apply all the low risk interventions are instituted for heavy or debilitated patients in a bed or wheel chair or on a foilet seat  Use restraints and bed monitors as ordered by the doctor  Allow the patient to ambulate only with assistance  Consider pask refacts of the medications that effects level of consciousness, gait and elimination when planning patient's care  Do not leave patients unattended in diagnostic or treatment areas  Accompany the patient while going to bathroom  Advice the patient to use grab bars near the tollet, bathtub, and shower  Make sure the family and other visitors understand the restrictions mentioned above  [Ight-risk interventions (45 or above)  Apply all the low and medium risk interventions  Tie red fall risk tag in the bed, wheel chair and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN	Remove excess equipment or furniture to make	a clear	4 4	. /-				1			
during procedure  Teach fall-prevention techniques, such as sitting up for a moment before rising from the bed  Bed wheels should be locked  Ensure that floor of the bathroom is dry and not slippery  Review medications for potential side effects that can promote falls  Use safety betts during movement in wheelchair  The patients are not ambulated by themselves. They are to be ambulated only with assistance  Meditum risk interventions (25-44)  Apply all the low risk interventions  Tie yellow fall risk tag in the bed and Wheel chair / Stretcher  Make sure that proper transfer precautions are instituted for heavy or debilitated patients in a bed or wheel chair or on a tollet seat  Use restraints and bed monitors as ordered by the doctor  Allow the patient to ambulate only with assistance  Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patients care.  Do not leave patients unattended in diagnostic or treatment areas  Accompany the patient while going to bathroom  Advice the patient to use grab bars near the tollat, bathtub, and shower  Make sure the family and other visitors understand the restrictions mentioned above  [Igh-Tisk Interventions (45 or above)  Apply all the low and medium risk interventions  Tie red fall risk tag in the bed, wheel chair and stretcher  Locate the high-Tisk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  Urinal/bedpan should be within easy reach (if appropriate)  Urinal/bedpan should be within easy reach (if appropriate)  If appropriate, consider using protection devices: satety belts  Signature & Emp. No. of RN	path		سرت ا	. •							
Teach fall-prevention techniques, such as sitting up for a moment before rising from the bed Bed wheels should be locked Encourage family participation in the patients care Ensure that floor of the bathroom is dry and not slippery Review medications for potential side effects that can promote fails Use safety belts during movement in wheelchair The patients are not ambulated by themselves. They are to be ambulated only with assistance    Make sure that proper transfer precautions are instituted for heavy or debilitated patients in a bed or wheel chair or an tollet seat Use restraints and bed monitors as ordered by the doctor Allow the patient to ambulate only with assistance   Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patient's care   Do not leave patients unattended in diagnostic or treatment areas   Do not leave patient use grab bars near the toilet, bathtub, and shower   Make sure the family and other visitors understand the restrictions mentioned above   Indinstructions (45 or a bove)	Keep the patient's bed in the low position at all times	sexcept		. /							
Bed wheels should be locked  Bed wheels should be locked  Ensure that floor of the bathroom is dry and not slippery  Review medications for potential side effects that can promote falls  Use safety belts during movement in wheelchair  The patients are not ambulated by themselves. They are to be ambulated only with assistance  (Actimur isk interventions (25-44)  Apply all the low risk interventions  Tie yellow fall risk tag in the bed and Wheel chair / Stretcher  Make sure that proper transfer precautions are instituted for heavy or debilitated patients in a bed or wheel chair or on a toilet seat  Use restraints and bed monitors as ordered by the doctor  Allow the patient to ambulate only with assistance  Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patients care  Do not leave patients unattended in diagnostic or treatment areas  Accompany the patient to use grab bars near the toilet, bathtub, and shower  Make sure the family and other visitors understand the restrictions mentioned above  Interventions (45orabove)  Apply all the low and medium risk interventions  Tiered fall risk tag in the bed, wheel chair and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call belis as quickly as possible  Provide a commode at bedside (if appropriate)  Urinal/bedpan should be within easy reach (if appropriate)  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN		. <b>.</b>	•		* *						
Bed wheels should be locked Encourage family participation in the patient's care Ensure that floor of the bathroom is dry and not slippery Review medications for potential side effects that can promote falls Use safety belts during movement in wheelchair The patients are not ambulated by themselves. They are to be ambulated only with assistance  Locate that proper transfer precautions are instituted for heavy or debilitated patients in a bed or wheel chair or on a toilet seat Use restraints and bed monitors as ordered by the doctor Allow the patient to ambulate only with assistance Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patient's care Do not leave patients unattended in diagnostic or treatment areas Accompany the patient while going to bathroom Advice the patient to use grab bars near the toilet, bathtub, and shower Make sure the family and other visitors understand the restrictions mentioned above  Interventions (45 or above) Apply all the low and medium risk interventions Tie red fall risk tag in the bed, wheel chair and stretcher Locate the high-risk patients in a room close to the nurses' station  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN  Signature & Emp. No. of RN		up for a	. /		-	l	-				
Encourage family participation in the patient's care Ensure that floor of the bathroom is dry and not slippery Review medications for potential side effects that can promote falls Use safety belts during movement in wheelchair The patients are not ambulated by themselves. They are to be ambulated only with assistance Meditum risk interventions (25-44) Apply all the low risk interventions Tile yellow fall risk tag in the bed and Wheel chair / Stretcher Make sure that proper transfer precautions are instituted for heavy or debilitated patients in a bed or wheel chair or on a toilet seat Use restraints and bed monitors as ordered by the doctor Allow the patient to ambulate only with assistance Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patient's care Do not leave patients unattended in diagnostic or treatment areas Accompany the patient while going to bathroom Advice the patient to use grab bars near the toilet, bathtub, and shower Make sure the family and other visitors understand the restrictions mentioned above Iligin-iisk interventions (45-or above) Apply all the low and meditum risk interventions Tile red fall risk tag in the bed, wheel chair and stretcher Locate the high-risk patients in a room close to the nurses' station Answer these patients call bells as quickly as possible Provide a commode at bedside (if appropriate) Urinal/bedpen should be within easy reach (if appropriate) If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN					, 						
Review medications for potential side effects that can promote falls  Use safety belts during movement in wheelchair  The patients are not ambulated by themselves. They are to be ambulated only with assistance Medium risk interventions (25-44)  Apply all the low risk interventions (25-44)  Apply all the low risk interventions are instituted for heavy or debilitated patients in a bed or wheel chair or an a toilet seat  Use restraints and bed monitors as ordered by the doctor  Allow the patient to ambulate only with assistance  Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patient's care  Do not leave patients unattended in diagnostic or treatment areas  Accompany the patient while going to balthroom  Advice the patient to use grab bars near the toilet, bathtub, and shower  Make sure the family and other visitors understand the restrictions mentioned above   IIGh-risk interventions (45 or above)  Apply all the low and medium risk interventions  Tiered fall risk tag in the bed, wheel chair and stretcher  Locate the high-risk patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belsts				•							
Review medications for potential side effects that can promote fails  Use safety belts during movement in wheelchair  The patients are not ambulated by themselves. They are to be ambulated only with assistance  Medium risk interventions  Tie yellow fall risk tag in the bed and Wheel chair / Stretcher  Make sure that proper transfer precautions are instituted for heavy or debilitated patients in a bed or wheel chair or on a toilet seat  Use restraints and bed monitors as ordered by the doctor  Allow the patient to ambulate only with assistance  Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patient's care  Do not leave patients unattended in diagnostic or treatment areas  Accompany the patient while going to bathroom  Advice the patient to use grab bars near the toilet, bathtub, and shower  Make sure the family and other visitors understand the restrictions mentioned above  Ight isk interventions (45 or above)  Apply all the low and medium risk interventions  Tie red fail risk tag in the bed, wheel chair and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (flappropriate)  Encourage family members or other visitors to stay with the appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN	• ., .				,						
Do not leave patients unattended in diagnostic or treatment areas  Accompany the patients to use grab bars near the toilet, bathtub, and shower  Make sure the family and other visitors understand the restrictions mentioned above  Millshrisk interventions (35 or above)  Apply all the low and medium is interventions  Tie yellow fall risk tag in the bed and Wheel chair / Stretcher  Make sure that proper transfer precautions are instituted for heavy or debilitated patients in a bed or wheel chair or on a toilet seat  Use restraints and bed monitors as ordered by the doctor Allow the patient to ambutate only with assistance  Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patients care  Do not leave patients unattended in diagnostic or treatment areas  Accompany the patient while going to bathroom  Advice the patient to use grab bars near the toilet, bathtub, and shower  Make sure the family and other visitors understand the restrictions mentioned above  Ilightrisk interventions (45 or above)  Apply all the low and medium isk interventions  Tie red fall risk tag in the bed, wheel chair and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (flappropriate)  Encourage family members or other visitors to stay with them  If a ppropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN											
Use safety belts during movement in wheelchair The patients are not ambulated by themselves. They are to be ambulated only with assistance  Medium risk interventions (25-44) Apply all the low risk interventions Tie yellow fall risk tag in the bed and Wheel chair / Stretcher Make sure that proper transfer precautions are instituted for heavy or debilitated patients in a bed or wheel chair or on a toilet seat Use restraints and bed monitors as ordered by the doctor Allow the patient to ambulate only with assistance Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patient's care Do not leave patients unattended in diagnostic or treatment areas Accompany the patient while going to bathroom Advice the patient to use grab bars near the toilet, bathtub, and shower Make sure the family and other visitors understand the restrictions mentioned above Make sure the family and other visitors understand the restrictions mentioned above Apply all the low and medium risk interventions Tie red fall risk tag in the bed, wheel chair and stretcher Locate the high-risk patients in a room close to the nurses' station Answer these patients call bells as quickly as possible Provide a commode at bedside (if appropriate) Encourage family members or other visitors to stay with them If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN	•	hat can							]		
The patients are not ambulated by themselves. They are to be ambulated only with assistance  Medium risk interventions (25-44)  Apply all the low risk interventions Tie yellow fall risk tag in the bed and Wheel chair / Stretcher  Make sure that proper transfer precautions are instituted for heavy or debilitated patients in a bed or wheel chair or on a toilet seat  Use restraints and bed monitors as ordered by the doctor Allow the patient to ambulate only with assistance  Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patients care  Do not leave patients unattended in diagnostic or treatment areas  Accompany the patient while going to bathroom  Advice the patient to use grab bars near the toilet, bathtub, and shower  Make sure the family and other visitors understand the restrictions mentioned above  Ifigh-risk interventions (45 or above)  Apply all the low and medium risk interventions  Tie red fall risk tag in the bed, wheel chair and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: salety belts  Signature & Emp. No. of RN	promote falls										
be ambulated only with assistance    Medium risk interventions (25-44)	Use safety belts during movement in wheelchair		•/-	$\mathcal{A}$			_		[		
Apply all the low risk interventions Tie yellow fall risk tag in the bed and Wheel chair / Stretcher Make sure that proper transfer precautions are instituted for heavy or debilitated patients in a bed or wheel chair or on a toilet seat Use restraints and bed monitors as ordered by the doctor Allow the patient to ambulate only with assistance Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patients care Do not leave patients unattended in diagnostic or treatment areas Accompany the patient thile going to bathroom Advice the patient to use grab bars near the toilet, bathtub, and shower Make sure the family and other visitors understand the restrictions mentioned above Iligh-risk interventions (45 or above) Apply all the low and medium risk interventions Tie red fall risk tag in the bed, wheel chair and stretcher Locate the high-risk patients call bells as quickly as possible Provide a commode at bedside (frappropriate) Urinal/bedpan should be within easy reach (if appropriate) Encourage family members or other visitors to stay with them If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN	The patients are not ambulated by themselves. The	ey are to		/							
Apply all the low risk interventions Tie yellow fall risk tag in the bed and Wheel chair / Stretcher Make sure that proper transfer precautions are instituted for heavy or debilitated patients in a bed or wheel chair or on a toilet seat Use restraints and bed monitors as ordered by the doctor Allow the patient to ambulate only with assistance Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patient's care Do not leave patients unattended in diagnostic or treatment areas Accompany the patient while going to bathroom Advice the patient to use grab bars near the toilet, bathtub, and shower Make sure the family and other visitors understand the restrictions mentioned above #ifgh-risk interventions (45 or above) Apply all the low and medium risk interventions Tie red fall risk tag in the bed, wheel chair and stretcher Locate the high-risk patients in a room close to the nurses' station Answer these patients call belis as quickly as possible Provide a commode at bedside (if appropriate) Urinal/bedpan should be within easy reach (if appropriate)  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN	be ambulated only with assistance						-		1		
Tie yellow fall risk tag in the bed and Wheel chair / Stretcher Make sure that proper transfer precautions are instituted for heavy or debilitated patients in a bed or wheel chair or on a toilet seat Use restraints and bed monitors as ordered by the doctor Allow the patient to ambulate only with assistance Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patient's care Do not leave patients unattended in diagnostic or treatment areas Accompany the patient while going to bathroom Advice the patient to use grab bars near the toilet, bathtub, and shower Make sure the family and other visitors understand the restrictions mentioned above ####################################	Medium risk interventions (25 - 44)		<del>-</del>			<del>                                     </del>	-	1	}		
Make sure that proper transfer precautions are instituted for heavy or debilitated patients in a bed or wheel chair or on a toilet seat  Use restraints and bed monitors as ordered by the doctor  Allow the patient to ambulate only with assistance  Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patient's care  Do not leave patients unattended in diagnostic or treatment areas  Accompany the patient while going to bathroom  Advice the patient to use grab bars near the toilet, bathtub, and shower  Make sure the family and other visitors understand the restrictions mentioned above  Igh-risk interventions (45 or above)  Apply all the low and medium risk interventions  Tie red fall risk tag in the bed, wheel chair and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN	Apply all the low risk interventions										
for heavy or debilitated patients in a bed or wheel chair or on a toilet seat  Use restraints and bed monitors as ordered by the doctor  Allow the patient to ambulate only with assistance  Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patient's care  Do not leave patients unattended in diagnostic or treatment areas  Accompany the patient while going to bathroom  Advice the patient to use grab bars near the toilet, bathtub, and shower  Make sure the family and other visitors understand the restrictions mentioned above ### in the bed, wheel chair and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  Urinal/bedpan should be within easy reach (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN	Tie yellow fall risk tag in the bed and Wheel chair/St	tretcher				}			_	ı,	
Use restraints and bed monitors as ordered by the doctor  Allow the patient to ambulate only with assistance  Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patient's care  Do not leave patients unattended in diagnostic or treatment areas  Accompany the patient while going to bathroom  Advice the patient to use grab bars near the toilet, bathtub, and shower  Make sure the family and other visitors understand the restrictions mentioned above  **Ifigh-risk interventions**(45 or above)**  Apply all the low and medium risk interventions  Tie red fall risk tag in the bed, wheel chair and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  Urinal/bedpan should be within easy reach (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN	Make sure that proper transfer precautions are in	stituted						-			
Use restraints and bed monitors as ordered by the doctor Allow the patient to ambulate only with assistance Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patient's care Do not leave patients unattended in diagnostic or treatment areas Accompany the patient while going to bathroom Advice the patient to use grab bars near the toilet, bathtub, and shower Make sure the family and other visitors understand the restrictions mentioned above Iligh-risk interventions (450r above) Apply all the low and medium risk interventions Tie red fall risk tag in the bed, wheel chair and stretcher Locate the high-risk patients in a room close to the nurses' station Answer these patients call bells as quickly as possible Provide a commode at bedside (if appropriate) Urinal/bedpan should be within easy reach (if appropriate) Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN	for heavy or debilitated patients in a bed or wheel	chair or						-			
Allow the patient to ambulate only with assistance  Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patient's care  Do not leave patients unattended in diagnostic or treatment areas  Accompany the patient while going to bathroom  Advice the patient to use grab bars near the toilet, bathtub, and shower  Make sure the family and other visitors understand the restrictions mentioned above  **Iigh-risk interventions** (45 or above)  Apply all the low and medium risk interventions  Tie red fall risk tag in the bed, wheel chair and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  Urinal/bedpan should be within easy reach (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN			<i></i>	,							
Consider peak effects of the medications that effects level of consciousness, gait and elimination when planning patients care  Do not leave patients unattended in diagnostic or treatment areas  Accompany the patient while going to bathroom  Advice the patient to use grab bars near the toilet, bathtub, and shower  Make sure the family and other visitors understand the restrictions mentioned above  **Iigh-risk interventions (45 or abovc)**  Apply all the low and medium risk interventions  Tie red fall risk tag in the bed, wheel chair and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  Linal/bedpan should be within easy reach (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN	Use restraints and bed monitors as ordered by the c	doctor									
of consciousness, gait and elimination when planning patient's care  Do not leave patients unattended in diagnostic or treatment areas  Accompany the patient while going to bathroom  Advice the patient to use grab bars near the toilet, bathtub, and shower  Make sure the family and other visitors understand the restrictions mentioned above High-risk interventions (45 or abovc)  Apply all the low and medium risk interventions  Tie red fall risk tag in the bed, wheel chair and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  Urinal/bedpan should be within easy reach (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN	Allow the patient to ambulate only with assistance										
patient's care  Do not leave patients unattended in diagnostic or treatment areas  Accompany the patient while going to bathroom  Advice the patient to use grab bars near the toilet, bathtub, and shower  Make sure the family and other visitors understand the restrictions mentioned above  iligh-risk interventions (45 or abovc)  Apply all the low and medium risk interventions  Tie red fall risk tag in the bed, wheel chair and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  Urinal/bedpan should be within easy reach (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN	•						-				
Do not leave patients unattended in diagnostic or treatment areas  Accompany the patient while going to bathroom  Advice the patient to use grab bars near the toilet, bathtub, and shower  Make sure the family and other visitors understand the restrictions mentioned above  **Iigh-risk interventions** (45 or abovc)  Apply all the low and medium risk interventions  Tie red fall risk tag in the bed, wheel chair and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  Urinal/bedpan should be within easy reach (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN	of consciousness, gait and elimination when p	lanning					1				
Accompany the patient while going to bathroom  Advice the patient to use grab bars near the toilet, bathtub, and shower  Make sure the family and other visitors understand the restrictions mentioned above  Iligh-risk interventions (45 or abovc)  Apply all the low and medium risk interventions  Tie red fall risk tag in the bed, wheel chair and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  Urinal/bedpan should be within easy reach (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN										٠.	
Accompany the patient while going to bathroom  Advice the patient to use grab bars near the toilet, bathtub, and shower  Make sure the family and other visitors understand the restrictions mentioned above  ilgh-risk interventions (45 or above)  Apply all the low and medium risk interventions  Tie red fall risk tag in the bed, wheel chair and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  Urinal/bedpan should be within easy reach (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN	Do not leave patients unattended in diagno	stic or	_				ļ		ļ		
Advice the patient to use grab bars near the toilet, bathtub, and shower  Make sure the family and other visitors understand the restrictions mentioned above  **Iigh-risk interventions (45 or abovc)**  Apply all the low and medium risk interventions  Tie red fall risk tag in the bed, wheel chair and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  Urinal/bedpan should be within easy reach (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN											
and shower  Make sure the family and other visitors understand the restrictions mentioned above  iligh-risk interventions (45 or abovc)  Apply all the low and medium risk interventions  Tie red fail risk tag in the bed, wheel chair and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  Urinal/bedpan should be within easy reach (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN			`			<u> </u>					
Make sure the family and other visitors understand the restrictions mentioned above  iigh-risk interventions (45 or above)  Apply all the low and medium risk interventions  Tie red fail risk tag in the bed, wheel chair and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  Urinal/bedpan should be within easy reach (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN	·	oathtub,									
restrictions mentioned above  iigh-risk interventions (45 or abovc)  Apply all the low and medium risk interventions  Tie red fall risk tag in the bed, wheel chair and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  Urinal/bedpan should be within easy reach (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN			<b>]</b>				ļ	ļ			
Apply all the low and medium risk interventions The red fall risk tag in the bed, wheel chair and stretcher Locate the high-risk patients in a room close to the nurses' station Answer these patients call bells as quickly as possible Provide a commode at bedside (if appropriate) Urinal/bedpan should be within easy reach (if appropriate) Encourage family members or other visitors to stay with them If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN	•	and the	,-	1				1			
Apply all the low and medium risk interventions  Tie red fall risk tag in the bed, wheel chair and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  Urinal/bedpan should be within easy reach (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN								ļ		Ì	
Tie red fall risk tag in the bed, wheel chair and stretcher  Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  Urinal/bedpan should be within easy reach (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN						-			-		
Locate the high-risk patients in a room close to the nurses' station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  Urinal/bedpan should be within easy reach (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN			<b> </b>			ļ		ļ. <u>.</u>	<del> </del>	=	
Station  Answer these patients call bells as quickly as possible  Provide a commode at bedside (if appropriate)  Urinal/bedpan should be within easy reach (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN			ļ					<del> </del>	<b></b>		
Answer these patients call bells as quickly as possible Provide a commode at bedside (if appropriate) Urinal/bedpan should be within easy reach (if appropriate) Encourage family members or other visitors to stay with them If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN	<u> </u>	nurses'		, ,	<i>'</i> .	ļ			-		
Provide a commode at bedside (if appropriate)  Urinal/bedpan should be within easy reach (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN		-1-	<del> </del>			<del>                                     </del>	-	<del> </del>	ļ	<del> </del>	
Urinal/bedpan should be within easy reach (if appropriate)  Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN		oie	<del> </del>			<u> </u>	<u> </u>	<del> </del>	<b></b>	<b> </b>	
Encourage family members or other visitors to stay with them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN	\		<del>                                     </del>			-	<del> </del>	<del>                                     </del>	<del>                                     </del>		
them  If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN	1					-	-	<del> </del>	<del> </del> -	<del> </del>	
If appropriate, consider using protection devices: safety belts  Signature & Emp. No. of RN		tay with		1	,	1					
Signature & Emp. No. of RN		u potetr	1	<del> </del>		<del> </del>	-	<del>                                     </del>	-		
Signature & Emp. No. of RN		. salety	,	];	<u>,</u> , •					1	
100			DN A	- July -	<del></del>	_	-	<u> </u>	<del>                                     </del>	<u> </u>	$\vdash$
Signature & Emp. No. of Sr. RN	Signature & Emp. No.	of RN	Thise	400		<u> </u>		<u> </u>		<u> </u>	
	Signature & Emp. No. of S	Sr. RN	Ma	Lan	Y		-				

vo

## MEDWAY HOSP ILS

### KODAMBAKKAM (HEART)

#9, 1st Main Road, United India Colony, Kodambakkam, Chennai,

Tamilnadu, India 044-2473 4455

care@medwayhospitals.com

Registration No

: MHI202481729

Patient Name

: SAMRUDDIN S M

Age

: 59

Gender

: Male

IP Number

: MMH/HM/IPH2024000093

Discharge Date

 $: 11/01/2024 \setminus 2:41:00PM$ 

Bill No

: MMH/HM/IPH202400088

Bill Date

: 11/01/2024

\1:39:37PM

Ward Name

: RADIAL LOUNGE

Bed Name

: RL-2

NO DUE



Approved By

Checked By